


Richard M. Marks, MD  
Professor and Director  
Division of Foot and Ankle  
Department of Orthopaedic Surgery  
Medical College of Wisconsin

## Adult Acquired Flat Foot

Adult Acquired Flatfoot also referred to as Pes Planovalgus is commonly caused by dysfunction of the Posterior Tibial Tendon. This tendon is primarily responsible for maintaining the arch as well as inverting the foot. The tendon typically fails as a result of degenerative changes and less commonly acute rupture. Risk factors for posterior tibial tendon dysfunction include obesity, age, hypertension, diabetes, female gender, and systemic inflammatory disorders such as rheumatoid arthritis. When the tendon is dysfunctional, patients may note collapse of their arch or the foot turning outward (abduction). As the deformity progresses, patients may suffer from lateral impingement symptoms or hindfoot arthritis.

If the posterior tibial tendon is actively inflamed, a course of immobilization with a boot or cast may be recommended. Other conservative treatment recommendations may include physical therapy, use of an anti-inflammatory agent, and/or orthotics. In advanced cases, bracing may be necessary. When conservative measures do not fail, surgical treatment may be considered.



Procedure In patients with a flexible deformity and minimal arthritic changes, surgery may entail debridement and repair to the dysfunctional tendon. In some cases, a posterior tibial tendon transfer may be warranted utilizing the flexor digitorum longus (FDL)

Following surgery you will be unable to place weight on your surgical extremity thus preoperative planning is essential. Prior to surgery, an appointment with a physical therapist will be made for instructional use of crutches or a walker as their use will be required postoperatively. The device will also be fitted to your height during this appointment. The crutches or walker will be issued at that appointment or arrangements will be made to obtain the device or arrangements will be through a medical supply company approved by your insurance. Some patients may opt to use a Rollator or wheel chair. These devices can be obtained through your local medical supply store. Please contact the office (414-805-7442) with the medical supply store of your choice and a prescription can be faxed in. Regardless of the modality used to maintain your weight bearing status, please practice in your home prior to surgery as repetition will reduce the risk of falls postoperatively. Removing throw rugs and clearing wider pathways through your home will also make navigating with crutches or walker easier and diminish the risk of falls.

During the period when strict elevation is required (the first ten days) you will need help with activities of daily living such as laundry, cooking, and cleaning. Please plan ahead and consider having friends or family stay with you. Driving is contraindicated during the acute postoperative recovery phase and may be prohibited for a longer period of time if your right foot requires immobilization. Showering will also be difficult during the recovery phase as you are unable to place weight on the surgical leg and cast/dressing needs to be kept clean and dry. Consider the use of a shower chair and/or hand held shower head. You will need to protect the leg by leaving it outside the shower as well as using bags or a plastic cast sleeve (brochure available in cast room) to ensure dressings remain dry.

## Postoperative Visits

### Day 0-10

Inpatient procedure: The procedure is performed on an outpatient basis.

Anesthesia: This is done under a general anesthetic.

Postoperative pain control

The use of a walker or crutches is required until you are able to place weight on your operative side.

xPain Control: Pain medications will be prescribed to be used as needed. Pre Operative nerve blocks can last between 8 to 12 hours; however, waiting to take pain medication until the block has completely worn off can result in increased breakthrough pain which can be difficult to manage. Please plan accordingly and take your medication promptly when sensation begins to return to the foot usually indicated by a tingling sensation in the toes or mild discomfort at the surgical site. Pain medications may be taken on a scheduled basis in the early postoperative recovery phase as this is when the pain is most intense.

### Day 10 –First PostOperative Visit

- x First postoperative visit with nonweightbearing x-rays, suture removal, application of cast.
- x Cast will remain on until week 6. This is necessary to allow for tendon and bony healing. Until then, cast must be kept clean, dry, and out of shower
- x Continue with nonweightbearing on operative leg and limited activities
- x May elevate extremity as needed

### Week 6

Cast removed, weightbearing x-rays obtained, advance weightbearing in a boot from 10-100% over the next 6-weeks with physical therapy with appropriate restrictions. Advancement of activities may be varied based on patients' past medical history and radiographic presentation.

### Week 12

Weaned from boot at about week 12 into regular shoes with orthotics. Increase activities as tolerated. It may require 10-12 months for complete rehabilitation. Compliance with home physical therapy protocol following discharge from formal physical therapy is key to improving strengthening and end-10(ke)-1181