Summer Programs Symposium

July 31st August 1st, 2024

Abstract Booklet

Summer Program for Undergraduate Research (SPUR)

Radhika Ramesh University of Wisconsin at Parkside
Kajana Satkunendrarajah, PhD
MCW Department of Neuroscience

Disruption of Forelimb Neural

Shreya Girish Case Western Reserve University
Kajana Satkunendrarajah, PhD
MCW Department of Neuroscience

The

Alexander Drobek Milwaukee School of Engineering
L. Tugan Muftuler, PhD

Candice Tacconi University of Lille, France
Stéphanie Olivier-Van Stichelen, PhD
MCW Department of Biochemistry

Hormone-related proteins are differentially *O*-GlcNAc-modified

4 in BeWo, and 5 in HEK293T cells. However, only 1 (ATXN2L) was detected in human placental total protein extracts.

Conclusion

This first verification validated the presence of 5 proteins in cell lines and 1 in human tissue sample: ATXN2L. The presence of the proteins in cell lines and not in human tissue is likely caused by the higher contamination of blood proteins, which led to loss of band definition on WB.

Protein Depletion Midi Spin from Thermo Scientific. This will remove proteins like albumin and transferrin so that we can produce an optimal image to visualize our proteins.

Keywords : O-GlcNAc; placenta; proteins; western-blot

Shayma Ouala Lille University; France.
Dawn Wenzel, PhD
MCW Department of Biochemistry

DISCOVERING NEW

Title: Decoding Sex-Specific Chronic Stress Effects on the Prelimbic Prefronta
Influence of REDD1/mTORC1 Signaling

Authors: A. Ramanujam[1, 2], V. Guda [2], M. K. Estes [2], B. Kurtoglu [2], D. Rodrigues de Ol
Mantsch

2,5-Dimethoxy-4-iodoamphetamine (DOI) regulates spine density and immediate early gene expression in the striatum

Peter Tagalakis[1], Jayashree Natajaran[2], Rutuja Borawke[2], John McCorvy[3], Coti Garcia Keller[2], Jennifer Tuscher[2]
1. University of Wisconsin-Parkside
2. MCW Department of Pharmacology and Toxicology
3. MCW Department of Cell Biology, Neurobiology, and Anatomy

Serotonergic psychedelics, including 2,5-Dimethoxy-4-iodoamphetamine (DOI), have garnered significant interest for their potential therapeutic benefit in treating substance use disorders, post-traumatic stress disorder, and depression[1,3,5,6,8]. DOI is one of several psychedelics that serves as an agonist for the 5-hydroxytryptamine receptor 2A ($5HTR_{2A}$). Although several recent studies have focused on psychedelic effects in the prefrontal cortex[2,4,7], less is known regarding the effects of psychedelics in striatal regions like the Nucleus Accumbens (NAc), which also expresses serotonin receptors, including $5HTR_{2A}$. Due to its association with motivation and positive and negative reinforcement, it is an excellent target for the treatment of depression and substance abuse. To test the hypothesis that DOI can similarly regulate spine density and plasticity-related gene expression in the striatum, we used a combination of *in vivo* and *in vitro* approaches. First, we administered 2 mg/kg DOI or saline vehicle to male and female rats intraperitoneally and monitored behavioral responses for 30 minutes. We found that 2mg/kg DOI significantly increased wet dog shake responses in female rats, a behavioral readout of serotonergic system activation in rodents hypothesized to parallel some

References

1. Canal, C. E., & Morgan, D. (2012). Head-twitch response in rodents induced by the hallucinogen 2,5-dimethoxy-4-iodoamphetamine: a comprehensive history, a re-evaluation of mechanisms, and its utility as a model. *Drug testing and analysis*, *4*(7-8), 556 576. https://doi.org/10.1002/dta.1333

2. Desouza, L. A., Benekareddy, M., Fanibunda, S. E., Mohammad, F., Janakiraman, B., Ghai, U., Gur, T., Blendy, J. A., & Vaidya, V. A. (2021). The Hallucinogenic Serotonin2A Receptor Agonist, 2,5-Dimethoxy-4-

Riya Tanwar University of Wisconsin-Madison
Gareth Pollin, PhD
Linda T. and John A. Mellowes Center for Genomic Sciences and Precision Medicine

Targeting PRMT5 Pathways to Disrupt Stress Granule Survival Mechanisms in Pancreatic Cancer

Background

Pancreatic cancer has a 5-year relative survival rate of less than 13%. The most common and

Ajay Katwala, *Marquette University*
Dr. Aron Geurts, PhD
MCW Department of Physiology

Heterotypic dual-vector ITRs increase efficiency in animal transgenesis

Background
Adeno-associated virus (AAV) is used to produce transgenic animal models to study human

capacity, but it has been shown to be overcome by a method utilizing a homotypic ITR dual-vector system, although with low efficiency. However, by utilizing a heterotypic ITR system compared to a homotypic ITR system, the low efficiency can be overcome and suggests a method of increasing transgenesis of large transgenes by utilizing a heterotopic

ITR vectors have greater efficiency at transducing HeLa cells. This could then be further applied to embryos, suggesting that transduction of larger transgenes can be done at a greater rate by utilizing a heterotypic ITR dual vector system with AAV.

Bao Ngoc Dinh Kenyon College
Daisy Sahoo, PhD,
MCW Department of Biochemistry

The Role of Modified HDL in Pro-atherogenic Signaling Cascades

Atherosclerosis, a major cause of cardiovascular disease, is a progressive disease that involves the buildup of fats, cholesterol and other substances on the artery walls (plaque). Plaques can cause arteries to narrow, blocking blood flow and increasing the likelihood of heart attack. High-density lipoproteins (HDL) play important roles in the pathogenesis of atherosclerosis due to its cardioprotective roles in removing peripheral cholesterol, along with its anti-inflammatory and antioxidant properties. Recent studies from our lab have demonstrated that HDL modified by reactive aldehydes has impaired cardioprotective properties in macrophages such as decreased macrophage migration, along with impaired ability to facilitate key steps in reverse cholesterol transport. Our goal in this study is to test the hypothesis that modification of HDL results in decreased macrophage migration. Specifically, we wanted to investigate the phosphorylation status of the MAP kinase pathways (ERK, JNK, p38) in macrophages treated with native HDL and modified HDL as these pathways are involved in macrophage migration. In order to do this, we modified HDL with malondialdehyde (MDA), and verified successful modification via immunoblot analyses. Next, we isolated peritoneal macrophages from wild-type (WT) and CD36 KO mice to compare the levels of phosphorylation of various signaling proteins of the MAP kinase pathways in the presence of native vs MDA-HDL. To do this, WT and CD36 KO mice were injected with thioglycolate and macrophages were harvested from peritoneal cavity, plated treated with native and MDA-HDL, and cells were lysed to collect cellular lysates. Preliminary data suggest that MDA-HDL delayed ERK 1/2 activation in WT macrophages while native HDL stimulates JNK in WT macrophages. Taken together, the data suggest that MDA-HDL may have a delayed activation period in comparison to native HDL in macrophages.

Key words: Cardiovascular disease, atherosclerosis, modified high-density lipoprotein (HDL), malondialehdyde (MDA), MAP kinase pathways

Alexandra Chafetz Kalamazoo College
Yiliang Chen, PhD
Department of Endocrinology and Molecular Medicine/Cardiovascular Center

---

Resistin Increases Cholesterol in Macrophages Contributing to Atherosclerosis

Background

Cardiovascular disease (CVD) remains the leading cause of death worldwide. Atherosclerosis, a condition characterized by the development of arterial plaques, is a primary contributor to heart attacks and strokes and serves as a key indicator of CVD progression. The pathogenesis of atherosclerosis involves various inflammatory pathways, with macrophages playing a crucial role in plaque formation. Macrophages are specialized immune cells that typically function to eliminate dead cells, debris, and metabolize substances like low-density lipoprotein (LDL) a form of cholesterol. These cells possess specific receptors that enable the recognition and uptake of cholesterol. One such receptor, cluster of differentiation 36 (CD36), recognizes oxidized LDL (oxLDL), a modified form of cholesterol resulting from the interaction between LDL and reactive oxygen species. While macrophages normally process oxLDL as part of cholesterol metabolism, excessive uptake can lead to their transformation into foam cells - swollen, dysfunctional cells that are a primary component of atherosclerotic plaques. This paper will examine the effects of resistin, a protein that appears to upregulate CD36 production, thereby potentially increasing foam cell formation and contributing to the progression of atherosclerosis.

Method

This experiment used mouse macrophages - Wild type (WT) and CD36 knock-out (CD36 KO). Macrophages

incubated with 1640 RPMI medium with fetal bovine serum. The cells were analyzed via flow cytometry and cholesterol assay to detect CD36 presence and cholesterol concentration, respectively.

Results

The results of the cholesterol assays indicated a (w)-422On(cat)7(ed)-157(a)12( )-158((w)-422On(cat)7(ed)-157(a)12( )-158((w)

Alyssa Bivens Wisconsin Lutheran College
Alexis Knotek
MCW Department of Microbiology and Immunology

Investigating MltG Regulation in *Enterococcus faecalis* Antibiotic Resistance

Background

Enterococci are opportunistic pathogens present within the human gut as commensal microbes. Their resistance to antibiotics allows them to grow when antibiotic treatments, for other illnesses, eliminate other microbes. This can result in lethal difficulties since their antibiotic resistance prevents most cell-wall antibiotic treatment from being effective. Enterococci can acquire resistance to vancomycin, an antibiotic that interferes with bacterial cell wall synthesis resulting in cell wall stress. This resistance to vancomycin is significantly reduced by deletion of MltG, a cell wall hydrolase predicted to be involved in peptidoglycan synthesis. The importance of MltG expression for vancomycin resistance prompted investigation into the regulation of MltG expression. Other enterococcal genes that

Felicity

Hannah Hotchkiss University of Wisconsin Oshkosh
Andrew Nencka, PhD
MCW Department of Radiology

Enhancing Large Language Models with Preference Expression: Weight Modification of the Gemma LLM for Improved AI Interaction and Bias Mitigation

Background

In the evolving field of artificial intelligence, enhancing the behavioral characteristics of large language models (LLMs) is critical for improving their human-like interaction. This study aimed to modify the weights of the Gemma LLM to develop and exhibit preferences. This enhancement is expected to lead to more nuanced and opinionated responses,

Morgan Kosch University of Wisconsin-Madison
Jessica Wagenknecht; Neshatul Haque, PhD; Michael T. Zimmermann, PhD
Linda T. and John A. Mellowes Center for Genomic Sciences and Precision Medicine at the Medical College of Wisconsin

Walker A and Walker B Interpretations of KRAS Protein Genetic Variants

Background:

The KRAS protein activates signal transduction pathways that lead to tumor growth. KRAS is the most

Jamal Beavers Iowa State University
Adam Greenberg, PhD
Medical College of Wisconsin Department of Biomedical Engineering

Post-Operative Cognitive Dysfunction and its Effects on Attentional Control

Introduction

Cognitive dysfunction following surgery is known to affect as much as 59% of the 60,000 patients undergoing surgical procedures that involve general anesthesia in the United States each day. This project aims to provide data that may ultimately allow us to develop auxiliary therapies that can help limit the effects of cognitive decline associated with general anesthesia. Post-operative cognitive dysfunction (POCD) is defined as an impairment of neural system functioning associated with the

may demonstrate that neuroinflammation is present in the limited brain regions associated with distractor filtering.

Emily Zhang Cornell University
Brandon Tefft, PhD
MCW-Marquette Department of Biomedical Engineering

RLD Peptide Enhances Endothelial Retention Under Shear Stress Conditions

Background

Establishing an endothelial monolayer can improve the patency of synthetic grafts (SGs) used in bypass surgeries. When endothelial cells (ECs) are exposed to shear stress induced by physiological blood flow, the loss of a functional endothelium can lead to a cascade of events including thrombogenesis, inflammation, and hyperplasia. Over the past decades, the RGD sequence has been intensively studied for its role in cell adhesion and its potential to help maintain endothelialization. While RGD has been found to promote greater cell retention, the

suitable alternative. In this study, we aim to analyze the RLD (Arg-Leu-Asp) peptide's ability to facilitate endothelial cell retention and compare its efficacy to RGD *in vitro*.

Methods

Human endothelial cells (HAECs) were cultured until 80% confluency. On day 1, pre-treated

HAECs were seeded in the flow path region using a rectangular gasket. After cells attached to the surface, fresh media was added to the dish to culture overnight. To assess the retention, shear stress was applied with a parallel plate flow chamber for five minutes. A flow rate of 3.18 mL/t4 591.46 Tm0 G[yBT/F2 11.04 Tfyu30 Tm( )] TJ.00000912 0 612 792 nBT2 reW*n.18 319.4> BDC609

Meg Dorrington Saint Louis University
Justin L. Grobe, PhD
MCW Departments of Physiology and Biomedical Engineering

Analysis

Courtney Griffin-Howard University
[Aprill Z. Dawson, PhD, MPH; Joni S. Willians, MD, MPH]
[MCW Center for Advancing Population Science]

Effective Clinical Trial

Zainab Khan

Vivien Blecking University of Wisconsin-Milwaukee

Olena Isaeva, PhD

MCW Department of Cell Biology, Neurobiology, and Anatomy

Juliet

Mae Jones University of Michigan
Elena Semina, PhD and Samuel Thompson, BS
MCW Department of Pediatrics, MCW Department of Ophthalmology and Visual Sciences

Investigating the Roles of ARHGAP18 and PRR12 in the Development of Microphthalmia

Background

Microphthalmia is an ocular condition where the eye is greater than two standard deviations smaller in

While phenotypes were present after breeding fish with variants in prr12a and arhgap18, no conclusions

Kayla Vanderhoef - University of Wisconsin- La Crosse
Ramani Ramchandran, PhD and Shubhangi Prabhudesai, PhD
MCW Department of Pediatrics, Research Institute

Fewer sources of platelet derived growth factor-